



FOXP2 gene

forkhead box P2

Normal Function

The *FOXP2* gene provides instructions for making a protein called forkhead box P2. This protein is a transcription factor, which means that it controls the activity of other genes. It attaches (binds) to the DNA of these genes through a region known as a forkhead domain. Researchers suspect that the forkhead box P2 protein may regulate hundreds of genes, although only some of its targets have been identified.

The forkhead box P2 protein is active in several tissues, including the brain, both before and after birth. Studies suggest that it plays important roles in brain development, including the growth of nerve cells (neurons) and the transmission of signals between them. It is also involved in synaptic plasticity, which is the ability of connections between neurons (synapses) to change and adapt to experience over time. Synaptic plasticity is necessary for learning and memory.

The forkhead box P2 protein appears to be essential for the normal development of speech and language. Researchers are working to identify the genes regulated by forkhead box P2 that are critical for learning these skills.

Health Conditions Related to Genetic Changes

FOXP2-related speech and language disorder


Several changes involving the *FOXP2* gene can result in *FOXP2*-related speech and language disorder, a condition that affects the development of speech and language starting in early childhood. Some affected individuals have a deletion that removes a small segment of chromosome 7, including the *FOXP2* gene and several neighboring genes. Other people with this condition have a mutation within the *FOXP2* gene itself. Less commonly, *FOXP2*-related speech and language disorder results from a rearrangement of the structure of chromosome 7 (such as a translocation) or from inheriting two copies of chromosome 7 from the mother instead of one from each parent (a phenomenon called maternal uniparental disomy or maternal UPD). It remains unclear how having two maternal copies of chromosome 7 affects the activity of the *FOXP2* gene.

The genetic changes that underlie *FOXP2*-related speech and language disorder disrupt the activity of the *FOXP2* gene. Because forkhead box P2 is a transcription factor, these changes affect the activity of other genes in the developing brain. Researchers are working to determine which of these genes are involved and how changes in their activity lead to abnormal speech and language development.

Chromosomal Location

Cytogenetic Location: 7q31.1, which is the long (q) arm of chromosome 7 at position 31.1

Molecular Location: base pairs 114,086,310 to 114,693,772 on chromosome 7 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- CAG repeat protein 44
- CAGH44
- forkhead/winged-helix transcription factor
- SPCH1
- TNRC10
- trinucleotide repeat containing 10

Additional Information & Resources

Educational Resources

- Coffee Break: Tutorials for NCBI Tools (2003): Talking About the Genetics of Talking
<https://www.ncbi.nlm.nih.gov/books/NBK2338/>
- Madame Curie Bioscience Database (2009): FOXP Genes, Neural Development, Speech and Language Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK7023/>

GeneReviews

- FOXP2-Related Speech and Language Disorders
<https://www.ncbi.nlm.nih.gov/books/NBK368474>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28FOXP2%5BTIAB%5D%29+OR+%28forkhead+box+P2%5BTIAB%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1080+days%22%5Bdp%5D>

OMIM

- FORKHEAD BOX P2
<http://omim.org/entry/605317>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_FOXP2.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=FOXP2%5Bgene%5D>
- HGNC Gene Family: Forkhead boxes
<http://www.genenames.org/cgi-bin/genefamilies/set/508>
- HGNC Gene Family: Trinucleotide repeat containing
<http://www.genenames.org/cgi-bin/genefamilies/set/775>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=13875
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/93986>
- UniProt
<http://www.uniprot.org/uniprot/O15409>

Sources for This Summary

- OMIM: FORKHEAD BOX P2
<http://omim.org/entry/605317>
- Feuk L, Kalervo A, Lipsanen-Nyman M, Skaug J, Nakabayashi K, Finucane B, Hartung D, Innes M, Kerem B, Nowaczyk MJ, Rivlin J, Roberts W, Senman L, Summers A, Szatmari P, Wong V, Vincent JB, Zeesman S, Osborne LR, Cardy JO, Kere J, Scherer SW, Hannula-Jouppi K. Absence of a paternally inherited FOXP2 gene in developmental verbal dyspraxia. *Am J Hum Genet.* 2006 Nov; 79(5):965-72. Epub 2006 Sep 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17033973>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1698557/>
- Fisher SE, Scharff C. FOXP2 as a molecular window into speech and language. *Trends Genet.* 2009 Apr;25(4):166-77. doi: 10.1016/j.tig.2009.03.002. Epub 2009 Mar 21. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19304338>

- Graham SA, Fisher SE. Understanding Language from a Genomic Perspective. *Annu Rev Genet.* 2015;49:131-60. doi: 10.1146/annurev-genet-120213-092236. Epub 2015 Oct 5.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/26442845>
- Lai CS, Fisher SE, Hurst JA, Vargha-Khadem F, Monaco AP. A forkhead-domain gene is mutated in a severe speech and language disorder. *Nature.* 2001 Oct 4;413(6855):519-23.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11586359>
- MacDermot KD, Bonora E, Sykes N, Coupe AM, Lai CS, Vernes SC, Vargha-Khadem F, McKenzie F, Smith RL, Monaco AP, Fisher SE. Identification of FOXP2 truncation as a novel cause of developmental speech and language deficits. *Am J Hum Genet.* 2005 Jun;76(6):1074-80. Epub 2005 Apr 22.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/15877281>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1196445/>
- Mueller KL, Murray JC, Michaelson JJ, Christiansen MH, Reilly S, Tomblin JB. Common Genetic Variants in FOXP2 Are Not Associated with Individual Differences in Language Development. *PLoS One.* 2016 Apr 11;11(4):e0152576. doi: 10.1371/journal.pone.0152576. eCollection 2016.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27064276>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4827837/>
- Vernes SC, Oliver PL, Spiteri E, Lockstone HE, Puliyadi R, Taylor JM, Ho J, Mombereau C, Brewer A, Lowy E, Nicod J, Groszer M, Baban D, Sahgal N, Cazier JB, Ragoussis J, Davies KE, Geschwind DH, Fisher SE. Foxp2 regulates gene networks implicated in neurite outgrowth in the developing brain. *PLoS Genet.* 2011 Jul;7(7):e1002145. doi: 10.1371/journal.pgen.1002145. Epub 2011 Jul 7.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21765815>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3131290/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/FOXP2>

Reviewed: September 2016
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services